


TOG-21-65A: VG-bearing Zone (350.6 to 352.25m)





General Description of Zone

A total of 30 clusters of visible gold were observed between 350.62 and 352.25m (red-outlined area, as shown above). Specks and blebs of gold vary in coarseness between <0.25 and 3 mm, mainly occurring tangentially to, or within, narrow white/translucent quartz stringers/patches of silicification. Narrow short micro-bands/wisps of VG get as coarse as 3mm long by 1mm wide.

The zone is characterized as a strongly (silica/albite/iron carbonate) altered felsite unit. Trace fine- to very fine-grained sphalerite is also present locally within a blebby white quartz-carbonate veinlet in this interval (at 351.6m), and is associated with small clots of dark green chlorite and trace fine-grained pyrite. Very fine- to fine-grained disseminated cubic pyrite is the dominant sulfide and occurs as scattered grains throughout the altered matrix and less commonly, within the odd white quartz-carbonate veinlet. No visible arsenopyrite was discerned during logging. Trace very fine- to fine-grained blebs of chalcopyrite not observed within VG-bearing interval, but occurs proximal to both the upper and lower contacts of the interval at 350.50m, 354.90m and 356.03m.

Close-up Photos of VG-bearing Zone





~352m: Fine-grained VG "floating" within clear/white quartz-carbonate veinlet. VG-bearing veinlet appears to crosscut narrow white quartz-carbonate extensional stringer with small clots of dark green chlorite.